

Correctional Facilities as Partners in Reducing HIV Disparities

Josiah D. Rich, MD, MPH,*†‡§ Ralph DiClemente, PhD,§ Judith Levy, PhD,|| Karen Lyda, LCSW, DNP,¶
Monica S. Ruiz, PhD, MPH,# David L. Rosen, PhD, MD,** and Dora Dumont, PhD, MPH,‡, On behalf of the
Centers for AIDS Research at the Social and Behavioral Sciences Research Network and the Centers for AIDS
Research–Collaboration on HIV in Corrections working group

Abstract: The United States now has the highest incarceration rate in the world. The majority of prison and jail inmates come from predominantly nonwhite and medically underserved communities. Although incarceration has adverse effects on both individual and community health, prisons and jails have also been used successfully as venues to provide health services to people with HIV who frequently lack stable health care. We review demographic trends shaping the difficulties in providing care to incarcerated people with HIV and recommend the Centers for AIDS Research Collaboration on HIV in Corrections as a model of interdisciplinary collaboration in addressing those difficulties.

Key Words: incarceration, prisons, underserved communities, co-occurring conditions

(*J Acquir Immune Defic Syndr* 2013;63:S49–S53)

From the *Brown University Medical School, Providence, RI; †Department of Medicine, Division of Infectious Diseases, The Miriam Hospital, Providence, RI; ‡The Center for Prisoner Health and Human Rights, Providence, RI; §Department of Behavioral Sciences, Rollins School of Public Health, Emory University, Atlanta, GA; ||University of Illinois at Chicago, Chicago, IL; ¶Division of Infectious Diseases, University of Colorado, School of Medicine, Denver, CO; #Department of Prevention and Community Health, George Washington University School of Public Health and Health Services, Washington, DC; and **Center for Infectious Diseases, University of North Carolina, Chapel Hill, NC.

Supported by the Centers for AIDS Research (grants P30AI027757, P30AI027763, P30AI036211, P30AI028697, P30AI060354, P30AI050409, P30AI036219, P30AI027742, P30AI027767, P30AI064518, P30AI051519, P30AI045008, P30AI087714, P30AI082151, P30AI042853, P30AI054999, P30AI078498, P30AI050410, P30AI073961, P30AI036214, and P30AI042845) and grants K24DA022112, K23DA029381, R01DA030762, R01DA18641, R01DA030778, R01DA027211, and R01DA027211 from the National Institute on Drug Abuse, National Institutes of Health and R01AA018944 from the National Institute on Alcohol Abuse and Alcoholism at the National Institutes of Health. Additional support for the supplement was provided by R13 MH-081733-01A1. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse or the National Institutes of Health.

Members of the Centers for AIDS Research at the Social and Behavioral Sciences Research Network and the Centers for AIDS Research–Collaboration on HIV in Corrections are listed in Appendix I.

The authors have no conflicts of interest to disclose.

Correspondence to: Josiah D. Rich, MD, MPH, Department of Medicine, Division of Infectious Diseases, The Miriam Hospital, 164 Summit Avenue, Providence, RI 02906 (e-mail: jrich@lifespan.org).

Copyright © 2013 by Lippincott Williams & Wilkins

INTRODUCTION

Two epidemics have converged in the United States over the past 30 years: mass incarceration and HIV. The United States now incarcerates a larger share of the population than any other nation in the world, and a larger percentage of our black population than South Africa did during apartheid.¹ Because incarceration is typically concentrated in low-income, medically underserved communities, many inmates have no contact with the health care system before incarceration and many of them are more appropriately candidates for mental illness or substance addiction treatment. Incarceration has significant adverse effects on both individual and community health. Paradoxically, though, jails (which typically hold those awaiting trial and those sentenced to less than 1 year) and prisons also create an opportunity to provide much-needed health services to the medically underserved. Correctional facilities have thus proved important sites for HIV diagnosis, treatment, and research.

Although the prevalence of HIV infection has declined recently among inmates,² it remains disproportionately higher than in the general population. Fourteen percent of all Americans infected with HIV, and 20% of black Americans infected with HIV, pass through the correctional system each year.³ Below, we identify several demographic trends and needs that in the absence of any fundamental change in national policy will continue to shape the potential for corrections-based research and treatment to reduce disparities in HIV.

Racial/Ethnic Disparities

People of color are disproportionately represented in the correctional population. Approximately 60% of inmates in state or federal prisons with sentences of over a year are black or Hispanic.⁴ According to the most recent data on the racial/ethnic characteristics of persons incarcerated in state and federal prisons, black non-Hispanic males had an imprisonment rate that was more than 6 times higher (3110 per 100,000 US residents) than the imprisonment rate for white non-Hispanic males (487 per 100,000) and almost 3 times higher than the rate for Hispanic males (1193 per 100,000).⁵ Black women are overrepresented in the US state and federal prison population. At year-end 2009, 1 in 703 black females was imprisoned compared with 1 in 1987 white females and 1 in 1356 Hispanic females.⁶

Communities of color also are disproportionately represented in the domestic HIV/AIDS epidemic. In 2009, blacks represented only 13% of the US population but accounted for 52% of all diagnoses of HIV/AIDS infection and a higher rate of HIV-related deaths (27.5 per 100,000) than any other racial or ethnic group.⁷ Similarly, Hispanics comprise 14% of the US population but have the second highest HIV prevalence rates in the nation after blacks.⁷ These racial/ethnic communities also represent the individuals most in need of health care coverage but least likely to have it. In turn, they have limited access to HIV care⁶ and other important health and social services, including addiction treatment, mental health services, and general primary care.⁸ These communities also experience greater health risks and disease burdens when compared with the general populations living in the same metropolitan area, county, or state.⁹

Women and Adolescents

The vast majority of inmates are adult males, but the proportions of female and adolescent inmates have grown more rapidly in the past 30 years, and both groups present distinct health issues. The health profile of women inmates (who constitute about 12% of jail and 7% of prison inmates) is substantially worse than that of men.¹⁰ Female inmates who have been tested have an HIV prevalence generally estimated at 1.9% versus 1.5% among male inmates. However, rates as high as 11.6% have been found among female inmates in New York.¹¹ Women inmates additionally exhibit high rates of mental illness, addiction, and histories of sexual abuse and trauma, all of which are associated with high-risk behaviors.^{12,13}

Like women, adolescents are a relatively small segment of the incarcerated population, but they are increasing more rapidly than adult males. In 2004, an estimated 800,000 people less than 20 years spent time in correctional or juvenile justice facilities.^{14,15} Incarcerated youth are disproportionately male, low income, and black or Hispanic. Incarcerated youth are more likely to report risky sexual behavior such as low prevalence of consistent condom use, multiple sex partners, and sex while high on drugs or alcohol. Similarly, HIV, sexually transmitted disease, and hepatitis C infection rates are higher than among other nonincarcerated youth.^{16,17} Finally, more than two-thirds of incarcerated youth have 1 or more psychiatric disorders.¹⁸ Rates of mild, moderate, and severe mental illness, from anxiety to posttraumatic stress disorders and psychosis, are generally higher among incarcerated compared with nonincarcerated youth.^{19,20} Additionally, being incarcerated may be a significant factor contributing to subsequent engagement in risk behaviors and development of adverse health outcomes.²¹ These excess rates of poor mental health in both women and adolescents complicate HIV treatment and prevention, and future research should emphasize integrated treatments tailored to the psychosocial needs of each group.

Aging Prisoners

Inmates more than 50 years constitute the fastest growing age group of prisoners in the United States. By the year 2030, about one-third of the US prison population is

expected to be 55 years of age or older.²² The “graying” of the US prison population is due to a combination of factors including stricter legislation that has led to longer sentences, mandatory minimum sentencing, and tighter parole criteria.²³ In addition, as is true of the general population, older adults who have been incarcerated are living longer either while in prison or after release. Few facilities offer specialized geriatric services.²⁴ Yet, acceleration in the rate of older inmates’ biological aging is well documented,²⁵ and prisoners more than 50 years are more likely than their younger counterparts to have multiple illnesses that require treatment.²⁶ These conditions, in turn, complicate the diagnosis and treatment of HIV along with its comorbidities.²⁷ These changing demographics have important implications for the structuring of prison health services including HIV prevention, treatment, and care, and for the lives of the prisoners themselves.

On release from incarceration, poor integration between the correctional and the public health systems can result in poor continuity of care for individuals, including older inmates, transitioning into the community.²⁸ Long-term prisoners who are released to the community as older adults face the challenges of reentry into a now unfamiliar society that likely has changed substantially from when they were first incarcerated.²⁹ Health care, including antiretroviral services for those who are HIV infected, needs to be accessed and undertaken within a new context and may differ by availability, provider, or regimen from what was available during incarceration. In addition, because both young and old former inmates typically are without immediate private or public health insurance on release, interruption of antiretroviral therapy (ART) can negatively affect their clinical well-being.^{30,31} Such findings point to the need to develop tailored discharge planning aimed at providing older inmates with structured plans, services, and skills to improve continuity of care for those living with HIV.

Comorbidities

Aging is also a factor in the co-occurrence of other medical conditions that impact patients with HIV. A study of older (more than 54 years) patients with HIV in an urban clinic found that 89% had at least 1 comorbid condition.³² However, comorbidities are not exclusive to older patients; a study of patients with HIV at a New York City clinic found that 92% had at least 1 comorbidity. In addition, only 84% self-reported their comorbid conditions, indicating that some clients may not have been aware of their chart-documented conditions.³³ Self-reporting was particularly low for obesity and hepatitis C. Hepatitis C is present in approximately 30% of all Americans with HIV,³⁴ but sensitivity of self-reported hepatitis C infection ranged from 66% to 77% in 3 studies.³³

Although there are limited data on coexisting medical conditions among prison and jail inmates with HIV, the rates of infectious and chronic conditions are significantly higher in correctional facilities than in the general population along with the prevalence of HIV.^{35–37} In particular, hepatitis C is present in 60%–90% of people who inject drugs,^{34,38} who are overrepresented among the correctional population. Because patients coinfect with both HIV and hepatitis C have also

been found to have more comorbidities than HIV mono-infected patients,³³ it is likely that inmates with HIV have a greater burden of coexisting diseases.

The potential for physical comorbidities is further heightened for the many inmates who have a mental illness or addiction in addition to HIV infection. Well over half of inmates at any given time have a DSM-IV (Diagnostic and Statistical Manual, Version IV) mental disorder and an estimated 16%–24% have a serious mental illness.¹² Estimates of the number of the incarcerated meeting DSM-IV criteria for drug dependence or abuse vary but remain well above 50%.¹² A comprehensive integrated approach to treating co-occurring conditions is critical, not only to forestall drug interactions but also to ensure maintenance of treatment after release from prison or jail. Such an approach is even more critical in the case of mental illness or addiction co-occurring with HIV because these comorbidities complicate treatments for HIV and hepatitis C that require a sustained adherence regime and careful monitoring for adverse drug interactions.³⁴

Linkage to Care

The number of AIDS-related deaths among prisoners has declined markedly, from 100 per 100,000 in 1995 to 9 per 100,000 in 2007.¹¹ These declines are largely attributable to the provision of medical care in prison, which includes access to life-sustaining ART. Although access to health care in prison can be challenging due to limited confidentiality, stigmatizing attitudes, and scarce resources, other attributes of incarceration such as the provision of basic necessities and diminished access to drugs, alcohol and tobacco can benefit prisoners' health.

Transitioning back into the community is difficult for many released prisoners. Released prisoners often resume lives that were chaotic before their imprisonment, in addition to which they must secure food, shelter, and employment; negotiate relationships strained by the prisoner's absence; confront issues of substance dependence and mental health; and abide by restrictions of parole and other legal sanctions.^{39,40} Not surprisingly, these issues often take precedence over efforts to establish health care in the community.⁴¹ Expedient resumption of health care also may be inhibited by the common state policy of disenrolling Medicaid recipients on incarceration.⁴² As a consequence, prisoners who received Medicaid before their imprisonment often lack this important source of health insurance on their release,⁴³ a time when they could arguably benefit from it the most. The Ryan White Care Act provides funding for medication and medical care for impoverished HIV-infected individuals, but a lag time between release from incarceration and program enrollment could similarly disrupt continuity of care and adherence to ART, though this is less likely than with Medicaid. It is unclear what will happen to the Ryan White Care act once the Affordable Care Act is implemented, but there is concern that shifting clients from Ryan White funding to Medicaid may cause additional disruption of care for patients transitioning from correctional facilities back to the community.

At the time of release, HIV-infected prisoners are typically provided a referral to a community physician and

a short supply of ART. Existing data suggest that these modest gestures are insufficient in bridging care from prison to the community. In perhaps the most dramatic documented example, less than 5% of HIV-infected inmates released from the Texas state prison system filled a free prescription for ART in time to avoid a disruption in medication,³⁰ and only 28% enrolled in a community HIV clinic within 90 days after their release.⁴⁴ The imperative to continue care, and in particular ART, among released prisoners is underscored by a recent multinational study demonstrating that initiation of ART among HIV-infected individuals resulted in a 96% reduction in transmission to their uninfected sexual partners.³¹ In addition to the deleterious (personal and public) health consequences that can result from lapses in care, these lapses also undermine the substantial financial investments by correctional systems in providing adequate HIV care for infected inmates.

Data exist on only a handful of programs to enhance continuity of care among released HIV-infected prisoners. These programs, which are largely based on a model of bridging case management, have demonstrated some ability to link released prisoners into care and to provide for other needs that are traditionally unmet.⁴⁵ However, none of the programs have yet demonstrated favorable outcomes for adherence or viral burden.^{46,47}

Addressing this need, several National Institutes of Health-funded projects are currently assessing promising strategies to enhance released prisoners' continuity of care and to reduce their infectivity. Among others, the projects include interventions using peer-based support, pharmacological treatment of substance dependence, patient-centered adherence counseling, transition case management services, and telemedicine; projects are also examining statewide patterns of linkage to care and the costs associated with linkage interventions. To avoid the myopic implementation of individual projects, investigators are coordinating survey instruments and outcomes across studies, so that the results will be mutually informative and have a stronger effect on HIV prevention and patient care.

Centers for AIDS Research–Collaboration on HIV in Corrections

The Centers for AIDS Research (CFARs) are funded by the National Institutes of Health to support and nurture translational research among institutions scattered across the nation. The 12 core facilities work together and with other institutions to provide expertise, resources, and services that have proved difficult to coordinate under more traditional funding streams. Recognizing the value of drawing correctional facilities into translational HIV research, the CFAR–Collaboration on HIV in Corrections (CFAR-CHIC) was formed in 2009. By facilitating collaboration among basic and clinical investigators and the criminal justice system, CFAR–Collaboration on HIV in Corrections can contribute to the development of new paradigms for the prevention, diagnosis, and treatment of HIV and associated conditions (including linkage to care after release) for individuals who are frequently medically complex and medically underserved.

CONCLUSIONS

Medical professionals must remain alert to the ethical dilemmas of working with institutions that contribute to health disparities by disproportionately incarcerating those who often more properly belong in treatment for mental illness or addiction. At the same time, partnerships between academic researchers and corrections have the potential to make inroads on the continuing disparities that exist in HIV diagnosis and treatment. Incarcerated individuals with HIV represent a medically complex and underserved population, and correctional facilities provide a means of reaching not only them but also the communities to which 95% of them will return. As those communities are frequently medically underserved as well, medical-correctional partnerships may prove a valuable contribution to the National HIV/AIDS Strategy of the United States, especially its goals of reducing HIV-related health disparities.

REFERENCES

- Alexander M. *The New Jim Crow: Mass Incarceration in the Age of Colorblindness*. New York, NY: The New Press; 2010.
- Spaulding AC, Seals RM, Page MJ, et al. HIV/AIDS among inmates and releasees from US correctional facilities, 2006: declining share of epidemic but persistent public health opportunity. *PLoS One*. 2009;4:e7558.
- Beckwith CG, Zaller ND, Fu JJ, et al. Opportunities to diagnose, treat, and prevent HIV in the criminal justice system. *J Acquir Immune Defic Syndr*. 2010;55(suppl 1):S49–S55.
- Harrison PM, Beck AJ. *Prisoners in 2005*. Washington, DC: U.S. Department of Justice, Office of Justice Programs, Bureau of Justice Statistics Bulletin; 2006. NCJ 215092. Available at: <http://www.ojp.usdoj.gov/bjs/pub/pdf/p05.pdf>. Accessed August 10, 2011.
- West HC, Sabol WJ, Greenman SJ. *Prisoners in 2009*. Washington, DC: U.S. Department of Justice, Office of Justice Programs, Bureau of Justice Statistics Bulletin; 2010. NCJ 231675. Available at: <http://bjs.ojp.usdoj.gov/content/pub/pdf/p09.pdf>. Accessed August 10, 2011.
- Kates J, Levi J. Insurance coverage and access to HIV testing and treatment: considerations for individuals at risk for infection and for those with undiagnosed infection. *Clin Infect Dis*. 2007;45(suppl 4):S255–S260.
- Centers for Disease Control and Prevention. *HIV/AIDS Surveillance Report 2009*. Vol. 21. Atlanta, GA: Centers for Disease Control and Prevention, Department of Health and Human Services; 2011. Available at: <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/>. Accessed August 10, 2011.
- Glaser JB, Greifinger RB. Correctional health care: a public health opportunity. *Ann Intern Med*. 1993;118(2):2139–2145.
- Liao Y, Bang D, Cosgrove S, et al. Surveillance of health status in minority communities—Racial and Ethnic Approaches to Community Health across the U.S. (REACH U.S.) Risk Factor Survey, United States, 2009. *MMWR Surveill Summ*. 2011;60:1–41. Available at: <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6006a1.htm>. Accessed August 10, 2011.
- Binswanger IA, Merrill JO, Krueger PM, et al. Gender differences in chronic medical, psychiatric, and substance-dependence disorders among jail inmates. *Am J Public Health*. 2010;100:476–482.
- Maruschak LM, Beavers R. *HIV in Prisons, 2007–08*. Washington, DC: US Department of Justice, Bureau of Justice Statistics; 2009. NCJ 228307.
- James DJ, Glaze LE. *Mental Health Problems of Prison and Jail Inmates*. Washington, DC: US Department of Justice, Bureau of Justice Statistics; 2006. Available at: <http://bjs.ojp.usdoj.gov/index.cfm?ty=pbdetail&iid=789>. Accessed August 10, 2011.
- Raj A, Rose J, Decker MR, et al. Prevalence and patterns of sexual assault across the life span among incarcerated women. *Violence Against Women*. 2008;14:528–541.
- Beck AJ, Harrison PM. *Prison and Jail Inmates at Midyear 2005*. Washington, DC: US Department of Justice, Bureau of Justice Statistics; 2006. NCJ 213133. Available at: <http://bjs.ojp.usdoj.gov/index.cfm?ty=pbdetail&iid=1007>. Accessed August 10, 2011.
- Sickmund M, Sladky A., Kang W. *Easy Access to Juvenile Court Statistics: 1985–2005*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention [distributor]; 2008. Available at: <http://www.ojjdp.ncjrs.gov/ojstatbb/ezajcs/>. Accessed August 10, 2011.
- Romero EG, Teplin LA, McClelland GM, et al. A longitudinal study of the prevalence, development, and persistence of HIV/sexually transmitted infection risk behaviors in delinquent youth: implications for health care in the community. *Pediatrics*. 2007;119:e1126–e1141.
- Templeton DJ. Sexually transmitted infection and blood-borne virus screening in juvenile correctional facilities: a review of the literature and recommendations for Australian centres. *J Clin Forensic Med*. 2006;13:30–36.
- Wasserman GA, McReynolds RS, Lucas CP, et al. The voice DISC-IV with incarcerated male youths: prevalence of disorder. *J Am Acad Child Adolesc Psychiatry*. 2002;41:314–321.
- Abram KM, Washburn JJ, Teplin LA. Posttraumatic stress disorder and psychiatric comorbidity among detained youths. *Psychiatr Serv*. 2007;58:1311–1316.
- Teplin LA, Abram KM, McClelland GM. Psychiatric disorders in youth in juvenile detention. *Arch Gen Psychiatry*. 2002;59:1133–1143.
- Valois RF, MacDonald JM, Bretous L, et al. Risk factors and behaviors associated with adolescent violence and aggression. *Am J Health Behav*. 2002;26:454–464.
- Enders SR, Paterniti DA., Meyers FJ. An approach to develop effective health care decision making for women in prison. *J Palliat Med*. 2005;8:432–439.
- Loeb SJ, Steffensmeier D, Myco PM. In their own words: older male prisoners' health beliefs and concerns for the future. *Geriatr Nurs*. 2007;28:319–329.
- Collins DR, Bird R. The penitentiary visit—a new role for geriatricians? *Age Ageing*. 2007;36:11–13.
- United Nations Office on Drugs and Crime. *Handbook on Prisoners with Special Needs*. New York, NY: United Nations; 2009.
- Aday RH. Golden years behind bars: special programs and facilities for elderly inmates. *Fed Probab*. 1994;58:47–54.
- Simone MJ, Appelbaum J. HIV in older adults. *Geriatrics*. 2008;63:6–12.
- Fazel S, Baillargeon J. The health of prisoners. *Lancet*. 2011;377:956–965.
- Williams BA, Lindquist K, Sudore RL, et al. Being old and doing time: functional impairment and adverse experiences of geriatric female prisoners. *J Am Geriatr Soc*. 2006;54:702–707.
- Baillargeon J, Giordano TP, Rich JD, et al. Accessing antiretroviral therapy following release from prison. *JAMA*. 2009;301:848–857.
- Shattock RJ, Warren M, McCormack S, et al. Turning the tide against HIV. *Science*. 2011;333:42–43.
- Shah SS, McGowan JP, Smith C, et al. Comorbid conditions, treatment, and health maintenance in older persons with human immunodeficiency virus in New York City. *Clin Infect Dis*. 2002;35:1238–1243.
- Weiss JJ, Osorio G, Ryan E, et al. Prevalence and patient awareness of medical comorbidities in an urban AIDS clinic. *AIDS Patient Care STDs*. 2010;24:39–48.
- Altice FL, Kamarulzaman A, Soriano VV, et al. Treatment of medical, psychiatric, and substance-use comorbidities in people infected with HIV who use drugs. *Lancet*. 2010;376:367–387.
- Harzke AJ, Baillargeon JG, Pruitt SL, et al. Prevalence of chronic medical conditions among inmates in the Texas prison system. *J Urban Health*. 2010;87:486–503. doi: 10.1007/s11524-010-9448-2.
- Binswanger IA, Krueger PM, Steiner JF. Prevalence of chronic medical conditions among jail and prison inmates in the USA compared with the general population. *J Epidemiol Community Health*. 2009;63:912–919.
- Wilper AP, Woolhandler S, Boyd JW, et al. The health and health care of US prisoners: results of a nationwide survey. *Am J Public Health*. 2009;99:666–672.
- Solomon L, Flynn C, Muck K, et al. Prevalence of HIV, syphilis, hepatitis B, and hepatitis C among entrants to Maryland correctional facilities. *J Urban Health*. 2004;81:25–37.
- Travis J. *But They All Come Back: Facing the Challenges of Prisoner Reentry*. Washington, DC: Urban Institute Press; 2005.
- Travis J, Solomon AL, Waul M. *From Prison to Home: The Dimensions and Consequences of Prisoner Reentry*. Washington, DC: The Urban

Institute; 2001. Available at www.urban.org/pdfs/from_prison_to_home.pdf. Accessed June 29, 2011.

41. Mallik-Kane K. *Returning Home. Illinois Policy Brief: Health and Prisoner Reentry*. Washington, DC: Urban Institute; 2005.
42. Morrissey JP, Dalton KM, Steadman HJ, et al. Assessing gaps between policy and practice in Medicaid disenrollment of jail detainees with severe mental illness. *Psychiatr Serv*. 2006;57:803–808.
43. Perez LM, Ro MJ, Treadwell HM. Vulnerable populations, prison, and federal and state Medicaid policies: avoiding the loss of a right to care. *J Correct Health Care*. 2009;15:142–149.
44. Baillargeon JG, Giordano TP, Harzke AJ, et al. Enrollment in outpatient care among newly released prison inmates with HIV infection. *Public Health Rep*. 2010;125(suppl 1):64–71.
45. Zaller ND, Holmes L, Dyl AC. Linkage to treatment and supportive services among HIV-positive ex-offenders in Project Bridge. *J Health Care Poor Underserved*. 2008;19:522–531.
46. Springer SA, Altice FL. Improving the care for HIV-infected prisoners: an integrated prison-release health model. In: Greifinger R, ed. *Public Health behind Bars: From Prisons to Communities*. New York, NY: Springer Science+Business Media; 2004:535–555.
47. Wohl DA, Scheyett A, Golin CE, et al. Intensive case management before and after prison release is no more effective than comprehensive pre-release discharge planning in linking HIV-infected prisoners to care: a randomized trial. *AIDS Behav*. 2011;15:356–364.

APPENDIX I

J. Adams (Duke), F. Aunon (Duke), A. Boutwell, MD (Partners Health Care), C. Carpenter, MD (Brown), H. Cassell, MPH (Vanderbilt), M. Comfort, PhD (RTI International), G. Culbert, BSN (University of Illinois at Chicago), N. DeSousa, MPH (Emory), T. Ghose, PhD (UPenn), T. Giordano, MD, MPH (Baylor College of Medicine), R. M. Grimes, PhD (Baylor/UT Houston), C. Haley (Vanderbilt), D. Huang, PhD (Rush), T. Kinlock, PhD (Friends Research), C. Krebs, PhD (RTI International), M. Malek, MD, MPH (LA Sheriff's Department), B. McGovern, MD (Tufts), J. Mitty, MD, MPH (Harvard), B. Montague, DO (Brown), J. Myers, PhD (University of California at San Francisco), A. Nijhawan, MD (Harvard), L. Ouellet, PhD (University of Illinois at Chicago), R. Pollini, PhD, MPH (Pacific Institute for Research and Evaluation), M. Ross, PhD, MD, MPH (Texas Medical Center), J. Schumacher, PhD (University of Alabama at Birmingham), P. Selwyn, MD (Montefiore), L. Solomon, PhD (Abt), D. Stone, MD (Tufts), L. Strick, MD, MS (Washington), L. E. Taylor, MD (Brown), M. Tolou-Shams, PhD (Brown), J. Tulskey, MD (University of California at San Francisco), S. Vermund, PhD (Vanderbilt), C. Wester, MD (Tennessee Department of Health), C. Williams, PhD (University of Illinois at Chicago), and S. Yard, MS (Washington).